

ONE REMARK ON CONSTRUCTION OF SEPARATED QUOTIENT-SPACE

YU.A. NERETIN

Datum 31 december 1995

ABSTRACT. We discuss elementary constructions of boundaries of symmetric spaces.

Let M be a compact metric space. Let $M = \bigcup_{\alpha \in A} M_\alpha$ be a partition of M ($M_\alpha \cap M_\beta = \emptyset$ if $\alpha \neq \beta$). Then the quotient-space A has canonical structure of a topological space. Recall that the set $P \subset A$ is closed if and only if $\bigcup_{\alpha \in P} M_\alpha$ is a closed subset in M . Let $a_1, a_2, \dots$ be a sequence in A . Let $a \in A$. Then $a_j \rightarrow a$ if there exist points $m_j \in M_{\alpha_j}, m \in M_\alpha$ such that $m_j \rightarrow m$ in M .

The space A is not need to be separated in Hausdorff sence. We are interested in the following question: how to construct separated analog of the quotientspace A ?

1. PRELIMINARIES. HAUSDORFF CONVERGENCE

Let $N \subset M$ be a closed subset. Denote by M_ε the set of all points $m \in M$ satisfying the condition: there exist $n \in N$ such that $\rho(m, n) < \varepsilon$. Let $[M]$ be the space of all closed subsets in M . Hausdorff distance $d(N, N')$ in $[M]$ between N and N' is the infimum of $\varepsilon > 0$ such that $N \subset N'_\varepsilon$ and $N'_\varepsilon \subset N$.

Recall that the metric space $[M]$ is compact. Recall also two simple facts on Hausdorff convergence. Denote by $\bar{S}$ the closure of the set S . Denote by $B_\varepsilon(m)$ the ball $\rho(m, n) < \varepsilon$.

Lemma 1. Let $N_j \in [M]$. Let K_σ ($\sigma \in \Sigma$) be all limit points of the sequence N_j . Then

- a) $\overline{\bigcup_{\sigma \in \Sigma} K_\sigma}$ coincides with the set of all $m \in M$ such that for all $\varepsilon > 0$ the set $N_j \cap B_\varepsilon(m)$ is nonempty for infinite number of j .
- b) $\bigcap_{\sigma \in \Sigma} K_\sigma$ coincides with the set of all $m \in M$ such that for all $\varepsilon > 0$ the set $N_j \cap B_\varepsilon(m)$ is nonempty for sufficiently large j .

2. CONSTRUCTION OF SEPARATED QUOTIENT-SPACE

Let a partition $M = \bigcup_{\alpha \in A} M_\alpha$ satisfies the following condition

*) for each $B \subset A$ the set $\overline{\bigcup_{\alpha \in B} M_\alpha}$ is the union of elements of the partition.

Fix an open subset $\mathcal{A} \subset A$ such that quotient-topology on $\mathcal{A}$ is separated. Denote by $\tilde{\mathcal{A}} \subset [M]$ the set of subsets $\overline{M_\alpha}$, $\alpha \in \mathcal{A}$. Let our data satisfy the condition

**) the map $\alpha \leftrightarrow \overline{M_\alpha}$ is a homeomorphism of the spaces $\mathcal{A}$ and $\tilde{\mathcal{A}}$.

Definition. The *separated quotient-space* $[[A]]$ is the closure of $\tilde{\mathcal{A}}$ in Hausdorff metrics.

Remark. Of course the construction depends on the set $\mathcal{A} \subset A$.

3. DESCRIPTION OF THE SET $[[A]]$

By lemma 1 and the condition *) the elements $N \in [[A]]$ are unions of elements M_α of the partition. Hence we associate to each $N \in [[A]]$ subset $S_N \subset A$ of all $\sigma \in A$ such that $M_\sigma \subset N$. Denote by $[A]$ the set of all subsets S_N . By construction we have canonical bijection $[[A]] \leftrightarrow [A]$.

The following proposition is evident.

Lemma 2. Let $S \subset A$. Then the following conditions are equivalent

- a) $S \in [A]$
- b) There exist a sequence $a \in A$ such that each limit point of a_j is an element of S and each element $s \in S$ is a limit of the sequence a_j in the quotient-topology on A .

Elements of $[A]$ we call *admissible subsets*.

4. EXAMPLE: COMPLETE COLLINEATIONS

Let M be the Grassmann manifold Gr_n of all n -dimensional subspaces in $\mathbb{C}^n \oplus \mathbb{C}^n$. Let $\lambda \in \mathbb{C}^* = \mathbb{C} \setminus 0$. Let $V \in Gr_n$. Define the subspace λV :

$$h \oplus p \in V \Leftrightarrow h \oplus \lambda p \in \lambda V$$

where $h \in \mathbb{C}^n \oplus 0$, $p \in 0 \oplus \mathbb{C}^n$. Consider the partition of Gr_n into $\mathbb{C}^*$ -orbits. Let $Op \subset Gr_n$ be the space of graphs of invertible operators. Of course the space Op coincide with the general linear group $GL_n(\mathbb{C})$. The quotient space $Op/\mathbb{C}^* = GL_n(\mathbb{C})/\mathbb{C}^*$ is the group $PGL_n(\mathbb{C})$ of invertible operators defined up to scalar multiplier.

We want to apply our construction to the space $M = Gr_n$ and $\mathcal{A} = PGL_n(\mathbb{C})$. We have to describe all admissible subsets in $Gr_n/\mathbb{C}^*$.

Example. Let $n = 2$. Consider the sequence $Q_n = \begin{pmatrix} 1 & 0 \\ 0 & n \end{pmatrix} \in PGL_2(\mathbb{C})$. Then the set of limits of Q_n in $Gr_2/\mathbb{C}^*$ consists of points $V_1, \dots, V_5$ (= subspaces in $\mathbb{C}^2 \oplus \mathbb{C}^2$) enumerated below:

$$\begin{aligned}
V_1 &: (x, y; 0, 0) \\
V_2 &: (x, y; 0, y) \\
V_3 &: (x, 0; 0, y) \\
V_4 &: (x, 0; x, y) \\
V_5 &: (0, 0; x, y)
\end{aligned}$$

where $x, y \in V$. The subspaces V_1, V_3, V_5 are stable points of the group $\mathbb{C}^*$. The $\mathbb{C}^*$ -orbits of V_2, V_4 are 1-dimensional complex curves.

Definition. Let $V \in Gr_n$. Then

- a) *Kernel* $\text{Ker } V = V \cap (\mathbb{C}^n \oplus 0)$
- b) *Image* $\text{Im } V$ is the projection of V to $0 \oplus \mathbb{C}^n$.
- c) *Domain* $\text{Dom } V$ is the projection of V to $\mathbb{C}^n \oplus 0$.
- d) *Indefiniteness* $\text{Indef } V = V \cap (0 \oplus \mathbb{C}^n)$.

Remark. Let $V \in Gr_n$. Then the subspace V induces by the obvious way the invertible operator

$$\text{Dom } V / \text{Ker } V \rightarrow \text{Im } V / \text{Indef } V$$

We denote this operator by $< V >$.

Definition. *Hinge* in $\mathbb{C}^n$ is a collection

$$\mathcal{P} = (Q_0, P_1, Q_1, P_2, Q_2, \dots, P_k, Q_k)$$

where Q_j, P_j are elements of Gr_n defined up to multiplier and 0.

$$\begin{aligned}
Q_j &= \text{Ker } Q_j \oplus \text{Indef } Q_j \\
P_j &\neq \text{Ker } P_j \oplus \text{Indef } P_j
\end{aligned}$$

1. For each $j = 1, 2, \dots, k$

$$\begin{aligned}
\text{Ker } P_j &= \text{Ker } Q_j = \text{Dom } P_{j+1} \\
\text{Im } P_j &= \text{Im } Q_j = \text{Indef } P_{j+1}
\end{aligned}$$

- 2.

$$\begin{aligned}
Q_0 &= \mathbb{C}^n \oplus 0 ; \text{Dom } P_1 = \mathbb{C}^n \\
Q_k &= 0 \oplus \mathbb{C}^n ; \text{Im } P_k = \mathbb{C}^n .
\end{aligned}$$

Remark. Let P be the graph of an invertible operator $\mathbb{C}^n \rightarrow \mathbb{C}^n$. Then

$$(\mathbb{C}^n \oplus 0, P, 0 \oplus \mathbb{C}^n)$$

is a hinge.

Remark. The elements $Q_0, \dots, Q_{k+1}$ of a hinge are completely defined by the elements $P_1, \dots, P_k$. The subspaces Q_j are fixed points of the group $\mathbb{C}^*$. The $\mathbb{C}^*$ -orbits of P_j are 1-dimensional complex curves.

Theorem. *The space $[PGL_n]$ of all admissible subsets in $Gr_n/\mathbb{C}^*$ coincides with the space of all hinges.*

The space $[PGL_n]$ coincide with the *complete collineation* space constructed by Semple (see [2]). It is a smooth algebraic variety and the group PGL_n is an open dense subset in $[PGL_n]$. On equivalence of these two constructions see [8]. Complete collineations is a partial case of complete symmetric varieties, see De Concini, Procesi [3].

5. EXAMPLE. FURSTENBERG-SATAKE COMPACTIFICATION OF RIEMANNIAN SYMMETRIC SPACE

We will only discuss the case $PGL_n(\mathbb{R})/SO(n)$. Consider the space $\mathbb{R}^n \oplus \mathbb{R}^n$ provided by a skew-symmetric bilinear form $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. Let $\mathcal{L}$ be the grassmannian of all Lagrangian subspaces in $\mathbb{R}^n \oplus \mathbb{R}^n$. Denote by $\mathbb{R}^*$ the multiplicative group of real positive numbers. This group acts on $\mathcal{L}$ by multiplications of linear relations on scalars.

Denote by R the open subset in $\mathcal{L}$ consisting of graphs of operators $S : \mathbb{R}^n \rightarrow \mathbb{R}^n$. It is easy to see that

$$\{\text{matrix } S \text{ is symmetric}\} \Leftrightarrow \{\text{the graph of } S \text{ is an element of } \mathcal{L}\}$$

The group $GL_n(\mathbb{R})$ acts on R by the formula $g : S \mapsto g^t S g$. The stabilizer of the point $S = E$ is the orthogonal group $O(n)$. Hence $GL_n(\mathbb{R})$ -orbit X of E is a homogeneous space $GL_n(\mathbb{R})/O(n)$. Points of X correspond to positive definite matrices S .

Now we apply the construction of the sections 2-3 to the space $\mathcal{L}$ and to the open subset $X = GL_n(\mathbb{R})/O(n)$. Then the completion consists of hinges

$$P = (Q_0, P_1, Q_1, \dots, P_k, Q_k)$$

such that $P_j \in \mathcal{L}, Q_j \in \mathcal{L}$ and the operators $\langle P_j \rangle$ (see section 4) are positive definite.

6. EXAMPLE. BOUNDARY OF BRUHAT-TITS BUILDING

Let $\mathbb{Q}_p$ be a p -adic field. Let M be the space of all $\mathbb{Z}_p$ -submodules in $\mathbb{Q}_p^n$. Let $B \subset M$ be the space of all lattices. The group $\mathbb{Q}_p^*$ act on M in a natural way. Then the corresponding separated quotient-space consists of collections

$$(R_0, T_1, R_1, \dots, T_k, R_k)$$

where $0 = R_0 \subset T_1 \subset R_1 \subset T_2 \dots \subset R_k = \mathbb{Q}_p^n$ are elements of M defined up to multiplier, R_j are subspaces and images of T_j in R_j/R_{j-1} are lattices.

I thanks C. De Concini, S.L.Tregub and E.B.Vinberg for discussion of this subject.

BIBLIOGRAPHY

1. E. Study, *Über die Geometrie der Kegelschnitte, insbesondere deren charakteristische Probleme*, Math. Ann., 27 (1886), 51-58.

2. I.G. Semple, *The variety whose points represent complete collineations of S_r on S'_r* , Rend. Math. 10 (1951), 201-280.
3. Fürstenberg, H., *A Poisson formula for semisimple Lie groups*, Ann. of Math. (2) 77 (1963), 335-386.
4. Satake, I., *On representations and compactifications of symmetric Riemannian spaces*, Ann. of Math. (2) 71 (1960), 77-110.
5. Alguneid, A.R., *Complete quadrics primals in four dimensional space*, Proc. Math. Phys. Soc. Egypt, 4 (1952), 93-104.
6. C. De Concini, C. Prochesi, *Complete symmetric varieties*, Lect. Notes Math., 996, 1-44.
7. Oshima, T., Sekiguchi, I., *Eigenspaces of invariant differential operators on an affine symmetric space*, Invent. Math. 57 (1980), 1-81.
8. Neretin, Yu. A., *On universal completions of complex classical groups*, Funct. Anal. Appl., 26:1.
9. Vinberg E.B., *On reductive algebraic semigroups*, Adv. Sov. Math., volume dedicated to E.B. Dynkin, to appear.
10. Neretin Yu.A., *Hinges and Study-Semple-Furstenberg-Satake-De Concini-Procesi- Oshima boundary*, to appear.

ADDRESS MOSCOW STATE INSTITUTE OF ELECTRONIC AND MATHEMATICS

Current address: Max-Planck-Institute für Mathematik, Bonn

E-mail address: neretin@mpim-bonn.mpg.de ; neretin@matan.miemstu.msk.su